Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	119,933	121,078	38,908	38,287	22,648	22,454
GEOGRAPHIC DIVISION						
Northeast	22,990	22,550	7,360	7,263	4,421	4,319
New England	6,615	6,598	2,210	2,190	1,413	1,273
Middle Atlantic	16,375	15,952	5,150	5,073	3,007	3,046
Midwest	26,248	26,283	7,873	7,839	4,533	4,738
East North Central	18,750	18,453	5,894	5,725	3,466	3,568
West North Central	7,497	7,830	1,978	2,114	1,068	1,170
South	41,393	42,916	12,637	12,753	7,228	7,114
South Atlantic	22,273	23,350	6,957	6,871	4,038	3,716
East South Central	6,726	6,826	1,907	1,977	1,076	1,239
West South Central	12,395	12,741	3,773	3,905	2,114	2,159
West	29,302	29,329	11,038	10,432	6,466	6,283
Mountain	9,208	9,109	3,469 ^a	2,947	2,037 ^a	1,653
Pacific	20,094	20,219	7,569	7,486	4,429	4,630
COUNTY TYPE						
Large Metro	66,195	65,728	21,977	21,427	12,801	12,572
Small Metro	36,587	38,181	11,858	11,965	6,827	7,007
250K – 1 Mil. Pop.	24,847	26,052	8,040	8,038	4,613	4,740
< 250K Pop.	11,739	12,129	3,818	3,927	2,214	2,267
Nonmetro	17,151	17,169	5,073	4,894	3,020	2,875
Urbanized	7,788	7,840	2,321	2,413	1,488	1,431
Less Urbanized	8,124	7,608	2,460	2,047	1,395	1,206
Completely Rural	1,239 ^a	1,721	293 ^a	434	137 ^a	238

Table 1.59AIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in
Thousands, 2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	47.3	47.0	15.3	14.9	8.9	8.7
GEOGRAPHIC DIVISION						
Northeast	49.4	48.1	15.8	15.5	9.5	9.2
New England	54.0	53.5	18.0	17.8	11.5	10.3
Middle Atlantic	47.8	46.1	15.0	14.7	8.8	8.8
Midwest	47.4	47.2	14.2	14.1	8.2	8.5
East North Central	48.7	47.7	15.3	14.8	9.0	9.2
West North Central	44.6	46.1	11.8	12.4	6.4	6.9
South	44.5	45.1	13.6	13.4	7.8	7.5
South Atlantic	45.5	46.5	14.2	13.7	8.2	7.4
East South Central	44.6	44.6	12.6	12.9	7.1	8.1
West South Central	42.9	42.9	13.1	13.2	7.3	7.3
West	49.9	49.0	18.8	17.4	11.0	10.5
Mountain	51.1	50.2	19.2 ^a	16.2	11.3 ^a	9.1
Pacific	49.3	48.5	18.6	18.0	10.9	11.1
COUNTY TYPE						
Large Metro	48.8	48.0	16.2	15.7	9.4	9.2
Small Metro	47.2	47.4	15.3	14.8	8.8	8.7
250K – 1 Mil. Pop.	48.3	47.8	15.6	14.7	9.0	8.7
< 250K Pop.	45.2	46.5	14.7	15.0	8.5	8.7
Nonmetro	42.4	42.7	12.5	12.2	7.5	7.2
Urbanized	45.6	46.7	13.6	14.4	8.7	8.5
Less Urbanized	41.3	39.7	12.5	10.7	7.1	6.3
Completely Rural	33.3 ^a	41.0	7.9	10.3	3.7	5.7

Table 1.59BIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages,
2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	6,278	6,373	4,745	4,735	2,468	2,522
GEOGRAPHIC DIVISION						
Northeast	1,073	1,110	840	854	479	442
New England	307	307	242	250	142	138
Middle Atlantic	767	803	598	605	337	304
Midwest	1,320	1,304	986	960	506	511
East North Central	933	910	693	663	364	368
West North Central	387	395	293	297	141	144
South	2,200	2,209	1,657	1,585	792	831
South Atlantic	1,185	1,087	894	781	418	393
East South Central	330	336	251	240	111	139
West South Central	686	786	512	563	264	298
West	1,684	1,749	1,261	1,336	690	738
Mountain	520	511	394	390	198	215
Pacific	1,163	1,238	867	946	492	523
COUNTY TYPE						
Large Metro	3,353	3,403	2,571	2,521	1,405	1,314
Small Metro	2,011	2,013	1,496	1,507	757 ^a	864
250K – 1 Mil. Pop.	1,393	1,398	1,030	1,076	544	600
< 250K Pop.	617	615	466	432	213	265
Nonmetro	914	957	677	707	306	344
Urbanized	354	445	264 ^a	351	116 ^a	174
Less Urbanized	490	423	364	293	167	141
Completely Rural	70	89	50	63	23	28

Table 1.60AIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in
Thousands, 2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	25.8	25.5	19.5	19.0	10.1	10.1
GEOGRAPHIC DIVISION						
Northeast	25.3	25.9	19.8	20.0	11.3	10.3
New England	28.1	27.6	22.1	22.4	13.0	12.4
Middle Atlantic	24.4	25.4	19.0	19.1	10.7	9.6
Midwest	24.7	24.0	18.5	17.6	9.5	9.4
East North Central	24.9	23.9	18.5	17.4	9.7	9.7
West North Central	24.3	24.1	18.4	18.2	8.9	8.8
South	24.6	23.9	18.5	17.1	8.8	9.0
South Atlantic	26.4	23.5	19.9 ^a	16.9	9.3	8.5
East South Central	22.8	22.8	17.4	16.2	7.7	9.4
West South Central	22.7	24.9	16.9	17.8	8.7	9.4
West	28.9	29.2	21.7	22.3	11.9	12.3
Mountain	29.1	27.8	22.0	21.2	11.1	11.7
Pacific	28.9	29.8	21.5	22.8	12.2	12.6
COUNTY TYPE						
Large Metro	25.7	25.6	19.7	18.9	10.8	9.9
Small Metro	26.7	25.7	19.9	19.2	10.1	11.0
250K – 1 Mil. Pop.	27.7	26.1	20.5	20.0	10.8	11.2
< 250K Pop.	24.8	24.9	18.7	17.5	8.5 ^a	10.7
Nonmetro	24.1	25.0	17.9	18.5	8.1	9.0
Urbanized	23.2	26.3	17.3	20.7	7.6	10.3
Less Urbanized	25.3	23.9	18.8	16.6	8.6	8.0
Completely Rural	22.1	24.9	15.7	17.6	7.3	7.8

 Table 1.60B
 Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	19,538	19,534	11,990	12,067	7,345	7,352
GEOGRAPHIC DIVISION						
Northeast	3,686	3,540	2,373	2,332	1,465	1,410
New England	1,112 ^a	1,007	778	701	509	446
Middle Atlantic	2,573	2,533	1,596	1,631	956	964
Midwest	4,285	4,168	2,574	2,537	1,555	1,513
East North Central	2,974	2,939	1,828	1,792	1,115	1,084
West North Central	1,312	1,229	746	745	440	429
South	6,780	6,999	4,022	4,094	2,402	2,442
South Atlantic	3,497 ^a	3,738	2,166	2,269	1,325	1,374
East South Central	1,049	1,091	570	623	319	376
West South Central	2,234	2,169	1,285	1,202	758	692
West	4,787	4,827	3,020	3,103	1,923	1,987
Mountain	1,448	1,498	880	885	533	534
Pacific	3,339	3,330	2,140	2,219	1,390	1,453
COUNTY TYPE						
Large Metro	10,405	10,500	6,587	6,708	4,030	4,179
Small Metro	6,536	6,417	3,949	3,867	2,443	2,318
250K – 1 Mil. Pop.	4,233	4,128	2,541	2,510	1,554	1,456
< 250K Pop.	2,303	2,289	1,408	1,357	890	862
Nonmetro	2,597	2,616	1,454	1,492	872	854
Urbanized	1,276	1,322	750	803	458	467
Less Urbanized	1,153	1,088	616	582	363	325
Completely Rural	168	206	88	107	51	62

Table 1.61AIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in
Thousands, 2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	57.3	56.9	35.2	35.2	21.6	21.4
GEOGRAPHIC DIVISION						
Northeast	61.0	57.8	39.3	38.1	24.2	23.0
New England	68.5	62.8	47.8	43.7	31.3	27.8
Middle Atlantic	58.2	56.1	36.1	36.1	21.6	21.3
Midwest	57.2	56.8	34.3	34.6	20.7	20.6
East North Central	57.9	58.0	35.6	35.4	21.7	21.4
West North Central	55.7	54.0	31.7	32.8	18.7	18.9
South	54.6	55.5	32.4	32.5	19.3	19.4
South Atlantic	55.4	58.0	34.3	35.2	21.0	21.3
East South Central	52.8	54.3	28.7	31.0	16.1	18.7
West South Central	54.3	52.1	31.2	28.9	18.4	16.6
West	59.0	58.6	37.2	37.7	23.7	24.1
Mountain	58.0	60.6	35.3	35.8	21.4	21.6
Pacific	59.5	57.8	38.1	38.5	24.8	25.2
COUNTY TYPE						
Large Metro	57.3	57.0	36.3	36.4	22.2	22.7
Small Metro	58.8	57.6	35.5	34.7	22.0	20.8
250K – 1 Mil. Pop.	60.3	58.3	36.2	35.4	22.1	20.6
< 250K Pop.	56.2	56.3	34.4	33.4	21.7	21.2
Nonmetro	54.2	55.1	30.3	31.4	18.2	18.0
Urbanized	60.4	59.9	35.5	36.4	21.7	21.2
Less Urbanized	49.5	52.0	26.4	27.8	15.6	15.5
Completely Rural	48.1	46.4	25.2	24.1	14.7	14.0

Table 1.61BIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2010
and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	94,117	95,171	22,174	21,485	12,835	12,580
GEOGRAPHIC DIVISION						
Northeast	18,231	17,900	4,147	4,077	2,476	2,467
New England	5,195	5,284	1,191	1,239	762	689
Middle Atlantic	13,036	12,616	2,956	2,838	1,714	1,778
Midwest	20,642	20,810	4,312	4,341	2,473	2,714
East North Central	14,843	14,605	3,373	3,270	1,986	2,117
West North Central	5,799	6,206	939	1,071	487	597
South	32,413	33,708	6,959	7,074	4,034	3,841
South Atlantic	17,591	18,525	3,897	3,820	2,296	1,949
East South Central	5,347	5,398	1,086	1,114	646	724
West South Central	9,475	9,786	1,976	2,139	1,092	1,168
West	22,831	22,752	6,756	5,993	3,852	3,557
Mountain	7,240	7,101	2,194 ^a	1,672	1,306 ^b	903
Pacific	15,591	15,652	4,562	4,321	2,546	2,654
COUNTY TYPE						
Large Metro	52,437	51,824	12,819	12,198	7,366	7,079
Small Metro	28,040	29,751	6,413	6,591	3,627	3,825
250K – 1 Mil. Pop.	19,221	20,526	4,469	4,453	2,515	2,684
< 250K Pop.	8,819	9,225	1,944	2,138	1,112	1,140
Nonmetro	13,641	13,596	2,942	2,696	1,842	1,677
Urbanized	6,158	6,073	1,307	1,259	914	789
Less Urbanized	6,481	6,096	1,480	1,172	866	740
Completely Rural	1,001 ^a	1,426	155 ^a	265	62 ^a	148

Table 1.62AIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in
Thousands, 2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	48.2	48.0	11.4	10.8	6.6	6.3
GEOGRAPHIC DIVISION						
Northeast	50.3	49.0	11.4	11.2	6.8	6.8
New England	54.5	55.0	12.5	12.9	8.0	7.2
Middle Atlantic	48.8	46.9	11.1	10.6	6.4	6.6
Midwest	48.6	48.5	10.1	10.1	5.8	6.3
East North Central	50.1	49.0	11.4	11.0	6.7	7.1
West North Central	45.1	47.4	7.3	8.2	3.8	4.6
South	45.3	46.0	9.7	9.6	5.6	5.2
South Atlantic	46.1	47.4	10.2	9.8	6.0	5.0
East South Central	45.9	45.7	9.3	9.4	5.5	6.1
West South Central	43.6	43.7	9.1	9.6	5.0	5.2
West	50.9	49.9	15.1	13.1	8.6	7.8
Mountain	52.7	51.3	16.0 ^a	12.1	9.5 ^b	6.5
Pacific	50.1	49.3	14.7	13.6	8.2	8.4
COUNTY TYPE						
Large Metro	50.2	49.3	12.3	11.6	7.0	6.7
Small Metro	47.7	48.3	10.9	10.7	6.2	6.2
250K – 1 Mil. Pop.	48.8	48.8	11.3	10.6	6.4	6.4
< 250K Pop.	45.5	47.2	10.0	10.9	5.7	5.8
Nonmetro	42.8	43.0	9.2	8.5	5.8	5.3
Urbanized	45.8	47.1	9.7	9.8	6.8	6.1
Less Urbanized	42.1	39.8	9.6	7.7	5.6	4.8
Completely Rural	32.8 ^a	41.9	5.1	7.8	2.0 ^a	4.3

Table 1.62BIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages,
2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	113,655	114,705	34,163	33,552	20,180	19,932
GEOGRAPHIC DIVISION						
Northeast	21,917	21,440	6,520	6,409	3,941	3,877
New England	6,308	6,291	1,968	1,941	1,271	1,135
Middle Atlantic	15,609	15,149	4,552	4,468	2,670	2,742
Midwest	24,927	24,978	6,886	6,878	4,028	4,227
East North Central	17,817	17,543	5,201	5,062	3,101	3,201
West North Central	7,111	7,435	1,685	1,817	927	1,026
South	39,193	40,707	10,980	11,168	6,436	6,283
South Atlantic	21,088	22,263	6,063	6,089	3,621	3,323
East South Central	6,396	6,489	1,656	1,737	965	1,100
West South Central	11,709	11,955	3,262	3,342	1,850	1,861
West	27,618	27,579	9,776	9,096	5,775	5,544
Mountain	8,688	8,598	3,075 ^a	2,557	1,839 ^a	1,438
Pacific	18,930	18,981	6,702	6,540	3,936	4,106
COUNTY TYPE						
Large Metro	62,842	62,324	19,406	18,906	11,395	11,258
Small Metro	34,576	36,168	10,362	10,458	6,070	6,143
250K – 1 Mil. Pop.	23,454	24,654	7,011	6,963	4,068	4,140
< 250K Pop.	11,122	11,514	3,352	3,495	2,002	2,003
Nonmetro	16,238	16,212	4,396	4,188	2,714	2,531
Urbanized	7,434	7,395	2,057	2,062	1,372	1,257
Less Urbanized	7,634	7,184	2,096	1,754	1,228	1,064
Completely Rural	1,169 ^a	1,633	243 ^a	372	114 ^a	210

Table 1.63AIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in
Thousands, 2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Geographic Characteristic	Lifetime (2010)	Lifetime (2011)	Past Year (2010)	Past Year (2011)	Past Month (2010)	Past Month (2011)
TOTAL	49.6	49.3	14.9	14.4	8.8	8.6
GEOGRAPHIC DIVISION						
Northeast	51.8	50.3	15.4	15.0	9.3	9.1
New England	56.5	56.1	17.6	17.3	11.4	10.1
Middle Atlantic	50.1	48.2	14.6	14.2	8.6	8.7
Midwest	49.8	49.7	13.8	13.7	8.1	8.4
East North Central	51.2	50.3	14.9	14.5	8.9	9.2
West North Central	46.7	48.4	11.1	11.8	6.1	6.7
South	46.7	47.4	13.1	13.0	7.7	7.3
South Atlantic	47.4	48.9	13.6	13.4	8.1	7.3
East South Central	46.9	46.9	12.1	12.6	7.1	8.0
West South Central	45.3	45.1	12.6	12.6	7.2	7.0
West	52.2	51.2	18.5	16.9	10.9	10.3
Mountain	53.5	52.7	18.9 ^a	15.7	11.3 ^a	8.8
Pacific	51.6	50.6	18.3	17.4	10.7	10.9
COUNTY TYPE						
Large Metro	51.2	50.5	15.8	15.3	9.3	9.1
Small Metro	49.4	49.7	14.8	14.4	8.7	8.4
250K – 1 Mil. Pop.	50.5	50.2	15.1	14.2	8.8	8.4
< 250K Pop.	47.4	48.7	14.3	14.8	8.5	8.5
Nonmetro	44.3	44.6	12.0	11.5	7.4	7.0
Urbanized	47.8	49.0	13.2	13.7	8.8	8.3
Less Urbanized	43.1	41.3	11.8	10.1	6.9	6.1
Completely Rural	34.3 ^a	42.5	7.1	9.7	3.3	5.5

Table 1.63BIllicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages,
2010 and 2011

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.